arXiv:math/0212317v2 [math.QA] 25 Dec 2002

SOLITONS, BOUNDARIES, AND QUANTUM AFFINE ALGEBRAS

GUSTAV W DELIUS

Abstract. This is a condensed write-up of a talk delivered at the Ramanujan International Symposium on Kac-Moody Lie algebras and Applications in Chennai in January 2002. The talk introduces special coideal subalgebras of quantum affine algebras which appear in physics when solitons are restricted to live on a half-line by an integrable boundary condition. We review how the quantum affine symmetry determines the soliton S-matrix in affine Toda field theory and then go on to use the unbroken coideal subalgebra on the half-line to determine the soliton reflection matrix. This gives a representation theoretic method for the solution of the reflection equation (boundary Yang-Baxter equation) by reducing it to a linear equation.

1. Introduction

Quantum affine algebras are quantum deformations of the universal enveloping algebras of affine Kac-Moody algebras, introduced by Drinfeld [1] and Jimbo [2]. They are therefore obviously related to the topic of this conference.

Solitons are localized finite-energy solutions of relativistic wave equations with special properties. In the quantum theory they lead to particle states. The relation between classical solitons and Kac-Moody algebras was stressed already by the Kyoto group [3]. More recently the role that quantum affine algebras play in the quantum theory of solitons was uncovered in [4].

By **Boundaries** I mean spatial boundaries. In this talk space will be taken to be one-dimensional, i.e., the real line. Introducing a boundary by imposing a boundary condition at some point restricts physics to the half-line.

In this talk we will knit these three topics together. We will do so by using a particularly rich family of quantum field theories. They are known as quantum affine Toda field theories and they posses all three: a quantum affine symmetry algebra [4], soliton solutions [7, 8], and integrable boundary conditions [15].

The main topic of the talk will be certain coideal subalgebras of quantum affine algebras that remain unbroken symmetry algebras after imposing an integrable boundary condition [30]. These algebras provide a new tool for the solution of the reflection equation.

Because the first part of the talk is simply an introduction to the known theory of solitons and quantum affine symmetry in affine Toda field theory, we will be very brief in the next two sections. We give references to the literature where more details can be found. These references are not always to the original works but often to more recent works because they may be more useful in the given context.

The transparencies from this talk are available on the web at
http://www.york.ac.uk/depts/maths/physics/delius/talks/iskmaa.pdf .

2. Solitons in Affine Toda field theory

2.1. **Field equations.** Associated to every affine Kac-Moody algebra $\hat{g}$ there is a relativistic field equation

$$\partial_x^2\phi - \partial_t^2\phi = \frac{1}{2i}\sum_{j=0}^{n} \eta_j \, \alpha_j^\vee \, e^{i(\alpha_j^\vee, \phi)} \tag{1}$$

where the field $\phi = \phi(x,t)$ takes values in the root space of the underlying finite dimensional Lie algebra g, $\alpha_0, \ldots, \alpha_n$ are the simple roots of $\hat{g}$ projected onto the root space of g, $(\ ,\ )$ is the Killing form, $\alpha_j^\vee = 2\alpha_j/(\alpha_j, \alpha_j)$ and the η_j are coprime integers such that $\sum_{j=0}^n \eta_j \alpha_j^\vee = 0$. The sine-Gordon model is the simplest example of an affine Toda field equation corresponding to $g = \hat{sl}_2$.

The field equations (1) are integrable in the sense that they posses an infinite number of integrals of motion in involution. This follows for example from the fact that they can be written as the zero-curvature condition for a gauge-connection taking its value in the affine Lie algebra $\hat{g}$ [5].

2.2. **Soliton solutions.** The Toda field equations (1) have degenerate vacuum solutions with $\phi = 2\pi\lambda$ for any fundamental weight λ of g. Thus there are kink solutions which interpolate between these vacua. They are stable due to their nonzero topological charge

$$T[\phi] = \phi(\infty) - \phi(-\infty) = 2\pi\lambda. \tag{2}$$



Furthermore there are solutions that describe an arbitrary number of kinks, each moving with its own velocity. Surprisingly, when two of these kinks meet they pass through each other and reemerge with their original shape restored. It is this property that shows that these kinks are solitons. An example of a two-soliton solution in the sine-Gordon model is shown on the right.

The topological charges of the fundamental solitons are weights in the fundamental representations of g. All solitons whose topological charge lie in the same fundamental representation have the same mass. A particle physicist would therefore say that they form a multiplet.

As an introduction to solitons from the particle physicsts viewpoint I recommend the book by Rajaraman [6]. A more classical presentation is given in [3]. For details on the solitons in affine Toda theory see [7, 8].

2.3. **Quantum solitons.** In the quantum theory we associate particle states with the soliton solutions. Let V_θ^μ be the space spanned by the solitons in multiplet μ with rapidity[1] θ.



Asymptotic two-soliton states span tensor product spaces $V_\theta^\mu \otimes V_{\theta'}^\nu$.

An incoming two-soliton state in $V_\theta^\mu \otimes V_{\theta'}^\nu$ with $\theta > \theta'$ will evolve during scattering into an outgoing state in $V_{\theta'}^\nu \otimes V_\theta^\mu$ with scattering amplitude given by the entry of the two-soliton S-matrix $S^{\mu\nu}(\theta - \theta')$. We represent this graphically as in the figure on the right.

---

[1] The rapidity θ parametrizes the energy and momentum of particles on the mass shell. If the mass is m then the energy is $E = m\cosh\theta$ and the momentum is $p = m\sinh\theta$.

2.4. **S-matrix factorization.** Due to integrability, the multi-soliton S-matrix factorizes into a product of two-soliton S-matrices. For the three-particle scattering process this is represented graphically as follows:



(3)

The compatibility of the two ways to factorize requires the S-matrix to be a solution of the Yang-Baxter equation

$$\begin{aligned}&(1\otimes S^{\mu\nu}(\theta-\theta'))(S^{\mu\lambda}(\theta-\theta'')\otimes 1)(1\otimes S^{\nu\lambda}(\theta'-\theta''))\\ &\qquad = (S^{\nu\lambda}(\theta'-\theta'')\otimes 1)(1\otimes S^{\mu\lambda}(\theta-\theta''))(S^{\mu\nu}(\theta-\theta')\otimes 1). \qquad (4)\end{aligned}$$

A classical reference on factorized S-matrices is [9].

Quantum affine algebras provide the key technique for finding solutions to the Yang-Baxter equation, as will be explained in the next section.

3. Quantum group symmetry

3.1. **Non-local symmetry charges.** Quantum affine Toda theory has symmetry charges $T_i, Q_i, \bar{Q}_i$, $i = 0,\dots,n$ which generate the $U_q(\hat{g})$ algebra with relations

$$\begin{gathered}[T_i, Q_j] = \alpha_i\cdot\alpha_j\, Q_j,\ \ [T_i,\bar{Q}_j] = -\alpha_i\cdot\alpha_j\, Q_j\\ Q_i\bar{Q}_j - q^{-\alpha_i\cdot\alpha_j}\bar{Q}_j Q_i = \delta_{ij}\,\frac{q^{2T_i}-1}{q_i^2-1},\end{gathered} \qquad (5)$$

where

$$q = e^{2\pi i\frac{1-\hbar}{\hbar}} \qquad (6)$$

and $q_i = q^{\alpha_i\cdot\alpha_i/2}$. They also satisfy the Serre relations. The charges $Q_i, \bar{Q}_i$ are non-local in the sense that they are obtained as space integrals of the time components of currents which are themselves non-local expressions. The details can be found in [4].

3.2. **Action on solitons.** Each soliton multiplet V_θ^μ carries a representation $\pi_\theta^\mu : U_q(\hat{g}) \to \mathrm{End}(V_\theta^\mu)$ of the quantum affine algebra. The symmetry acts on the multi-soliton states through the coproduct $\Delta : U_q(\hat{g}) \to U_q(\hat{g})\otimes U_q(\hat{g})$.

$$\begin{aligned}\Delta(Q_i) &= Q_i\otimes 1 + q^{T_i}\otimes Q_i,\\ \Delta(\overline{Q}_i) &= \overline{Q}_i\otimes 1 + q^{T_i}\otimes \overline{Q}_i,\\ \Delta(T_i) &= T_i\otimes 1 + 1\otimes T_i.\end{aligned} \qquad (7)$$

An explanation of why non-local symmetry charges have such a non-cocommutative coproduct can be found in [10].

3.3. **S-matrix as intertwiner.** The S-matrix has to commute with the action of any symmetry charge $Q \in U_q(\hat{g})$,

$$\begin{tikzcd} V_\theta^\mu\otimes V_{\theta'}^\nu \arrow[r, "(\pi_\theta^\mu\otimes\pi_{\theta'}^\nu)(\Delta(Q))"] \arrow[d, "S^{\mu\nu}(\theta-\theta')"] & V_\theta^\mu\otimes V_{\theta'}^\nu \arrow[d, "S^{\mu\nu}(\theta-\theta')"] \\ V_{\theta'}^\nu\otimes V_\theta^\mu \arrow[r, "(\pi_{\theta'}^\nu\otimes\pi_\theta^\mu)(\Delta(Q))"] & V_{\theta'}^\nu\otimes V_\theta^\mu \end{tikzcd} \qquad (8)$$

This determines the S-matrix uniquely up to an overall factor. Jimbo solved this intertwining equation for the vector representations of all quantum affine algebras in [2]. A practical technique for solving the intertwining equations in a large number of cases is the tensor product graph method, see e.g. [11].

The scalar prefactor which is not fixed by the Yang-Baxter equation can be determined by imposing other physical requirements such as unitarity, crossing symmetry and closure of the bootstrap. This has been done for a number of affine Toda theories, see e.g., [12, 13, 14]

3.4. **Yang-Baxter equation from Schur's lemma.** Because the tensor product modules are irreducible for generic rapidities, Schur's lemma implies that the following diagram is commutative up to a scalar factor

$$\begin{tikzcd}
V^\mu_\theta \otimes V^\nu_{\theta'} \otimes V^\lambda_{\theta''} \arrow[r, "S^{\mu\nu}(\theta-\theta')\otimes\mathrm{id}"] \arrow[d, "\mathrm{id}\otimes S^{\nu\lambda}(\theta'-\theta'')"] & V^\nu_{\theta'} \otimes V^\mu_\theta \otimes V^\lambda_{\theta''} \arrow[d, "id\otimes S^{\mu\lambda}(\theta-\theta'')"'] \\
V^\mu_\theta \otimes V^\lambda_{\theta''} \otimes V^\nu_{\theta'} \arrow[d, "S^{\mu\lambda}(\theta-\theta'')\otimes\mathrm{id}"] & V^\nu_{\theta'} \otimes V^\lambda_{\theta''} \otimes V^\mu_\theta \arrow[d, "S^{\nu\lambda}(\theta'-\theta'')\otimes\mathrm{id}"'] \\
V^\lambda_{\theta''} \otimes V^\mu_\theta \otimes V^\nu_{\theta'} \arrow[r, "id\otimes S^{\mu\nu}(\theta-\theta')"] & V^\lambda_{\theta''} \otimes V^\nu_{\theta'} \otimes V^\mu_\theta
\end{tikzcd} \tag{9}$$

Thus any S-matrix which satisfies the intertwining property (8) automatically satisfies the Yang-Baxter equation (at least up to a scalar factor).

4. Soliton reflection

4.1. **Integrable boundary conditions.** We now restrict the theory to the left half-line by imposing that at the boundary

$$\partial_x\phi = i\sum_{j=0}^{n}\epsilon_j\alpha_j^\vee \exp\left(\frac{i}{2}(\alpha_j^\vee,\phi)\right) \tag{10}$$

where the ϵ_j are free parameters. These boundary conditions were proposed by Corrigan et.al. because they preserve integrability for certain choices of the parameters [15]. There are other integrable boundary conditions [16, 17] and it will be interesting to extend the observations from the next section to these.

4.2. **Soliton reflection.** For $a_n^{(1)}$ affine Toda theory the soliton solutions which satisfy the boundary conditions (10) were determined in [18] by a nonlinear method of images. These solutions describe an incoming soliton with rapidity θ being converted into an outgoing anti-soliton with rapidity $-\theta$. In the quantum theory such reflection processes are described by a reflection matrix $K^\mu(\theta)$ which maps from the incoming multiplet V^μ_θ to the outgoing conjugate multiplet $V^{\bar\mu}_{-\theta}$. The entries of the reflection matrix are the reflection amplitudes.



4.3. **Factorization of reflection matrices.** If the boundary condition preserves integrability then the multi-soliton reflection matrices have to factorize.



(11)

The compatibility condition between the two ways of factorizing the two-soliton reflection matrix is the *reflection equation* [19]

$$\begin{aligned}(1\otimes K^{\nu}(\theta'))S^{\nu\bar{\mu}}(\theta+\theta')(1\otimes K^{\mu}(\theta))S^{\mu\nu}(\theta-\theta')&\\ = S^{\bar{\nu}\bar{\mu}}(\theta-\theta')(1\otimes K^{\mu}(\theta))S^{\mu\bar{\nu}}(\theta+\theta')(1\otimes K^{\nu}(\theta')).&\quad (12)\end{aligned}$$

This equation is also sometimes known as the *boundary Yang-Baxter equation.* Like the Yang-Baxter equation (4) it is a non-linear functional matrix equation. It was solved to find the reflection matrix for the sine-Gordon solitons in [20] and then for the vector solitons in $a_n^{(1)}$ affine Toda theory in [21].

Solutions of the reflection equation are not only needed to describe the reflection of particles off a boundary. They are also used to construct integrable quantum spin chains with a boundary and they provide the boundary Bolzmann weights for solvable models of statistical mechanics. Unfortunately the reflection equation has proven difficult to solve directly except for matrices of very small dimension [22, 23], for diagonal matrices [24, 25] and a few other cases [26, 21, 27]. The two known systematic techniques for the construction of reflection matrices are the fusion procedure [28] and the extended Hecke algebraic techniques [29] but these have so far only produced a small subset of the possible solutions.

A way to find solutions to the reflection equation with the help of quantum affine algebras has only been discovered very recently [30]. This method arose by studying the non-local symmetries of affine Toda theory on the half-line, as will be described in the next section.

5. Boundary quantum group symmetry

5.1. **Symmetry charges.** In the presence of the boundary the non-local charges of the bulk theory are no longer conserved. Rather we found [30] that the boundary condition (10) breaks the symmetry to a subalgebra $\mathcal{B}_\epsilon \subset U_q(\hat{g})$ generated by

$$\widehat{Q}_i = Q_i + \bar{Q}_i + \hat{\epsilon}_i q^{T_i}, \ i = 0, \dots, n. \qquad (13)$$

Note that the symmetry algebra depends on the boundary parameters $\hat{\epsilon}_i$ (which are related to the ϵ_i in (10) by a rescaling). We derived these conserved charges using first order boundary conformal perturbation theory but we believe the result to hold exactly. For $\hat{g} = a_n^{(1)}$ we were able to show that this is the case, at least on shell, as explained in the next section. We can now report that recent calculations performed in collaboration with Alan George establish the symmetry also for $\hat{g} = d_n^{(1)}$. It is the subject of ongoing work to extend this to all affine Toda field theories with an integrable boundary condition.

5.2. **Reflection Matrix as Intertwiner.** The reflection matrix has to commute with the action of any symmetry charge $\hat{Q} \in \mathcal{B}_\epsilon \subset U_q(\hat{g})$, i.e., the following diagram is commutative:

$$\begin{tikzcd}
V_\theta^\mu \arrow[r, "\pi_\theta^\mu(\hat{Q})"] \arrow[d, "K^\mu(\theta)"] & V_\theta^\mu \arrow[d, "K^\mu(\theta)"] \\
V_{-\theta}^{\bar{\mu}} \arrow[r, "\pi_{-\theta}^{\bar{\mu}}(\hat{Q})"] & V_{-\theta}^{\bar{\mu}}
\end{tikzcd} \tag{14}$$

If the symmetry algebra $\mathcal{B}_\epsilon$ is "large enough" so that V_θ^μ and $V_{-\theta}^{\bar{\mu}}$ are irreducible modules of $\mathcal{B}_\epsilon$, then this linear equation determines the reflection matrix uniquely up to an overall factor. As we will see below, this is the case for example for the vector representation of $a_n^{(1)}$.

5.3. **Coideal property.** The residual symmetry algebra $\mathcal{B}_\epsilon$ turns out not to be a Hopf algebra because it does not have a coproduct $\Delta : \mathcal{B}_\epsilon \to \mathcal{B}_\epsilon \otimes \mathcal{B}_\epsilon$. Using the coproduct (7) of $U_q(\hat{g})$ we find

$$\Delta(\hat{Q}_i) = (Q_i + \bar{Q}_i) \otimes 1 + q^{T_i} \otimes \hat{Q}_i. \tag{15}$$

Thus the symmetry algebra $\mathcal{B}_\epsilon$ is a left coideal of $U_q(\hat{g})$ in the sense that

$$\Delta(\hat{Q}) \in U_q(\hat{g}) \otimes \mathcal{B}_\epsilon \quad \text{for all } \hat{Q} \in \mathcal{B}_\epsilon. \tag{16}$$

This allows it to act on multi-soliton states in the presence of a boundary because the coproduct can be used to define the action of $\mathcal{B}_\epsilon$ on any tensor product of a $U_q(\hat{g})$-module (solitons) with a $\mathcal{B}_\epsilon$-module (boundary states). We would like to stress that this is the first time that a symmetry algebra in physics has been found that is not a bialgebra.

5.4. **The Reflection Equation from Schur's lemma.** If $\mathcal{B}_\epsilon$ is "large enough" so that the tensor product modules are irreducible, then by Schur's lemma the following diagram is commutative up to a scalar factor

$$\begin{tikzcd}
V_\theta^\mu \otimes V_{\theta'}^\nu \arrow[r, "\mathrm{id} \otimes K^\nu(\theta')"] \arrow[d, "S^{\mu\nu}(\theta-\theta')"] & V_\theta^\mu \otimes V_{-\theta'}^{\bar{\nu}} \arrow[d, "S^{\mu\bar{\nu}}(\theta+\theta')"'] \\
V_{\theta'}^\nu \otimes V_\theta^\mu \arrow[d, "\mathrm{id} \otimes K^\mu(\theta)"] & V_{-\theta'}^{\bar{\nu}} \otimes V_\theta^\mu \arrow[d, "\mathrm{id} \otimes K^\mu(\theta)"] \\
V_{\theta'}^\nu \otimes V_{-\theta}^{\bar{\mu}} \arrow[d, "S^{\nu\bar{\mu}}(\theta+\theta')"] & V_{-\theta'}^{\bar{\nu}} \otimes V_{-\theta}^{\bar{\mu}} \arrow[d, "S^{\bar{\nu}\bar{\mu}}(\theta-\theta')"'] \\
V_{-\theta}^{\bar{\mu}} \otimes V_{\theta'}^\nu \arrow[r, "\mathrm{id} \otimes K^\nu(\theta')"] & V_{-\theta}^{\bar{\mu}} \otimes V_{-\theta'}^{\bar{\nu}}
\end{tikzcd} \tag{17}$$

Thus any reflection matrix which satisfies the intertwining property (14) automatically satisfies the reflection equation (12) (at least up to a scalar factor).

5.5. **Calculating Reflection Matrices.** Solving the quantum group intertwining property (14) is much easier than solving the reflection equation (12) directly because the intertwining property is a linear equation.

To illustrate this new technique we present here the derivation of the reflection matrix for the vector solitons in $a_n^{(1)}$ Toda theory. Using the representation matrices

$$\pi_\theta^\mu(\hat{Q}_i) = x\, e^{i+1}{}_i + x^{-1}\, e^i{}_{i+1} + \hat{\epsilon}_i \left((q^{-1}-1)\, e^i{}_i + (q-1)\, e^{i+1}{}_{i+1} + \mathbf{1}\right) \tag{18}$$

the intertwining property (14) gives the following set of linear equations for the entries of the reflection matrix:

$$0 = \hat{\epsilon}_i (q^{-1} - q) K^i{}_i + x\, K^i{}_{i+1} - x^{-1}\, K^{i+1}{}_i, \tag{19}$$

$$0 = K^{i+1}{}_{i+1} - K^i{}_i, \tag{20}$$

$$0 = \hat{\epsilon}_i\, q\, K^i{}_j + x^{-1}\, K^{i+1}{}_j,\ j \neq i, i+1, \tag{21}$$

$$0 = \hat{\epsilon}_i\, q^{-1}\, K^j{}_i + x\, K^j{}_{i+1},\ j \neq i, i+1. \tag{22}$$

If all $|\hat{\epsilon}_i| = 1$ then one finds the solution

$$K^i{}_i(\theta) = \left(q^{-1} \, (-q\, x)^{(n+1)/2} - \hat{\epsilon}\, q\, (-q\, x)^{-(n+1)/2} \right) \frac{k(\theta)}{q^{-1} - q}, \tag{23}$$

$$K^i{}_j(\theta) = \hat{\epsilon}_i \cdots \hat{\epsilon}_{j-1}\, (-q\, x)^{i-j+(n+1)/2}\, k(\theta),\ \text{for } j > i, \tag{24}$$

$$K^j{}_i(\theta) = \hat{\epsilon}_i \cdots \hat{\epsilon}_{j-1} \hat{\epsilon}\, (-q\, x)^{j-i-(n+1)/2}\, k(\theta),\ \text{for } j > i, \tag{25}$$

that is unique up to an overall numerical factor $k(\theta)$. If all $\hat{\epsilon}_i = 0$ then the solution is diagonal. For other values for the $\hat{\epsilon}_i$ there are no solutions. This is in agreement with the restrictions on the boundary parameters found by imposing classical integrability [15].

We have applied the same technique to derive the reflection matrices for any representation of $\hat{sl}_2$ [31]. We are currently calculating the reflection matrices for the vector representation for all families of affine Lie algebras, mirroring the work done by Jimbo for R-matrices [2].

5.6. **Boundary Bound States.** Particles can bind to the boundary, creating multiplets of boundary bound states. These span finite dimensional representations $V^{[\lambda]}$ of the symmetry algebra $\mathcal{B}_\epsilon$. The reflection of particles off boundary bound states is described by intertwiners

$$K^{\mu[\lambda]}(\theta) : V^\mu_\theta \otimes V^{[\lambda]} \to V^{\bar{\mu}}_{-\theta} \otimes V^{[\lambda]}. \tag{26}$$

Unfortunately nothing is known so far about the representation theory of the algebras $\mathcal{B}_\epsilon$. One objective of this talk is to encourage research into this direction. Ideally one would like to be able to calculate the branching rules for representations of $U_q(\hat{g})$ into irreducible representations of $\mathcal{B}_\epsilon$.

6. Reconstruction of symmetry

In this last section I want to show how to reconstruct the symmetry algebra of a theory with boundary from the knowledge of at least one of its reflection matrices.

Let us assume that for one particular $U_q(\hat{g})$ representation V^μ_θ we know the reflection matrix $K^\mu(\theta) : V^\mu_\theta \to V^{\bar{\mu}}_{-\theta}$. We define the corresponding $U_q(\hat{g})$-valued L-operators in terms of the universal R-matrix $\mathcal{R}$ of $U_q(\hat{g})$,

$$L^\mu_\theta = (\pi^\mu_\theta \otimes \text{id})\, (\mathcal{R}) \in \text{End}(V^\mu_\theta) \otimes U_q(\hat{g}), \tag{27}$$

$$\bar{L}^{\bar{\mu}}_\theta = \left(\pi^{\bar{\mu}}_{-\theta} \otimes \text{id}\right) (\mathcal{R}^{\text{op}}) \in \text{End}(V^{\bar{\mu}}_{-\theta}) \otimes U_q(\hat{g}). \tag{28}$$

Here $\mathcal{R}^{\text{op}}$ is the opposite universal R-matrix obtained by interchanging the two tensor factors. From these L-operators we construct the matrices

$$B^\mu_\theta = \bar{L}^{\bar{\mu}}_\theta \, (K^\mu(\theta) \otimes 1)\, L^\mu_\theta \in \text{End}(V^\mu_\theta, V^{\bar{\mu}}_{-\theta}) \otimes U_q(\hat{g}). \tag{29}$$

In index notation this becomes

$$(B^\mu_\theta)^\alpha{}_\beta = (\bar{L}^{\bar{\mu}}_\theta)^\alpha{}_\gamma (K^\mu(\theta))^\gamma{}_\delta (L^\mu_\theta)^\delta{}_\beta \in U_q(\hat{g}). \tag{30}$$

We find that for all θ the $(B_\theta^\mu)^\alpha{}_\beta$ are elements of a coideal subalgebra $\mathcal{B}$ which commutes with the reflection matrices. It is easy to check the coideal property:

$$\Delta\left((B_\theta^\mu)^\alpha{}_\beta\right) = (\bar{L}_\theta^{\bar\mu})^\alpha{}_\delta (L_\theta^\mu)^\sigma{}_\beta \otimes (B_\theta^\mu)^\delta{}_\sigma. \tag{31}$$

Also any $K^\nu(\theta') : V_{\theta'}^\nu \to V_{-\theta'}^{\bar\nu}$ which satisfies the appropriate reflection equation commutes with the action of the elements $(B_\theta^\mu)^\alpha{}_\beta$

$$K^\nu(\theta') \circ \pi_{\theta'}^\nu((B_\theta^\mu)^\alpha{}_\beta) = \pi_{-\theta'}^{\bar\nu}((B_\theta^\mu)^\alpha{}_\beta) \circ K^\nu(\theta'), \tag{32}$$

Applying the above construction to the vector solitons in affine Toda theory and expanding in powers of $x = e^\theta$ gives

$$B_\theta^\mu = B + x\sum_{l=0}^{n}(q^{-1}-q)\, e^{l+1}{}_l \otimes \left(Q_l + \bar{Q}_l + \hat\epsilon_l\, q^{T_l}\right) + \mathcal{O}(x^2). \tag{33}$$

This shows that the charges were correct to all orders.

The B-matrices can be shown to satisfy the quadratic relations

$$\check{P}R^{\bar\nu\bar\mu}(\theta-\theta')\check{P}\,\overset{1}{B_\theta^\mu}\,R^{\mu\bar\nu}(\theta+\theta')\,\overset{2}{B_{\theta'}^\nu} = \overset{2}{B_{\theta'}^\nu}\,\check{P}R^{\nu\bar\mu}(\theta+\theta')\check{P}\,\overset{1}{B_\theta^\mu}\,R^{\mu\nu}(\theta-\theta'), \tag{34}$$

Thus they generate a *reflection equation algebra* in the sense of Sklyanin [32].

7. Summary and Discussion

In the first half of the talk we reviewed how quantum affine algebras are used to find the solutions of the Yang-Baxter equation that describe soliton scattering in affine Toda theory. In the second half we then described the recently discovered analogous technique for soliton reflection off integrable boundaries. The main points were the following

- A boundary condition breaks the symmetry to a subalgebra $\mathcal{B}_\epsilon$ of the quantum affie algebra $U_q(\hat g)$, depending on the boundary parameters ϵ_i.
- The residal symmetry algebra $\mathcal{B}_\epsilon$ is not a Hopf algebra but only a coideal subalgebra of $U_q(\hat g)$.
- The symmetry implies that the reflection matrices are intertwiners of $\mathcal{B}_\epsilon$ representations. Because for generic rapidity the $U_q(\hat g)$ modules V_θ^μ spanned by the solitons are irreducible also as modules of the subalgebra $\mathcal{B}_\epsilon$, the intertwining property determines the reflection matrices uniquely up to a scalar factor.
- Because also the tensor product modules $V_\theta^\mu \otimes V_{\theta'}^\nu$ are irreducible as $\mathcal{B}_\epsilon$ modules, Schur's lemma implies that the reflection matrices that satisfy the intertwining property are also automatically solutions of the reflection equation.
- The intertwining property leads to a set of linear equations for the entries of the reflection matrix which is relatively straightforward to solve. We have thus obtained a very practical method of finding solutions to the reflection equation.
- At special imaginary values of the rapidity θ a soliton multiplets $V_t^\mu heta$ can become reducible and boundary bound states can form, spanning irreducible representations of $\mathcal{B}_\epsilon$. Thus the physically interesting study of boundary bound states is connected to the study of the branching rules from $U_q(\hat g)$ to $\mathcal{B}_\epsilon$.

While in this talk we concentrated on affine Toda field theories, a similar story unfolds in the principal chiral models. There the theory in the bulk has a Yangian symmetry, which in the presence of a boundary is broken to a twisted Yangian. This was discovered in [33] and was used to find reflection matrices in [33, 34].

We found the boundary quantum groups $\mathcal{B}_\epsilon$ by studying concrete boundary quantum field theories. However the question can also be approached purely mathematially. One would like to classify and construct all coideal subalgebras of quantum affine algebras or Yangians that qualify as boundary quantum groups in the sense that their intertwiners are solutions to the boundary Yang-Baxter equation. This would lead to a classification of solutions of the boundary Yang-Baxter equation in terms of boundary quantum groups in the same spirit as the classification of solutions of the Yang-Baxter equation in terms of representations of quantum groups.

I hope to have given you in this talk a glimpse of a whole new field of enquiry related to Kac-Moody algebras with applications in both physics and mathematics.

Note added:

Since this contribution was written in May 2002 several interesting papers have appeared on the subject of reflection matrices and coideal subalgebras. Examples include [35, 37, 36]

Acknowledgments

The work presented here was performed in collaboration with Niall MacKay and our graduate students Ben Short and Alan George. My work is supported by an EPSRC advanced fellowship.

References

[1] V. G. Drinfeld, "Quantum groups," *J. Sov. Math.* **41** (1988) 898–915.

[2] M. Jimbo, "Quantum R matrix for the generalized Toda system," *Commun. Math. Phys.* **102** (1986) 537–547.

[3] T. Miwa, M. Jimbo, and E. Date, *Solitons*. Cambridge University Press, Cambridge, 2000. Differential equations, symmetries and infinite-dimensional algebras, Translated from the 1993 Japanese original by Miles Reid.

[4] D. Bernard and A. Leclair, "Quantum group symmetries and nonlocal currents in 2-D QFT," *Commun. Math. Phys.* **142** (1991) 99–138.

[5] D. I. Olive and N. Turok, "Local conserved densities and zero curvature conditions for Toda lattice field theories," *Nucl. Phys.* **B257** (1985) 277.

[6] R. Rajaraman, "Solitons and instantons. an introduction to solitons and instantons in quantum field theory,". Amsterdam, Netherlands: North-Holland (1982) 409p.

[7] T. J. Hollowood, "Solitons in affine Toda field theories," *Nucl. Phys.* **B384** (1992) 523–540.

[8] D. I. Olive, N. Turok, and J. W. R. Underwood, "Affine Toda solitons and vertex operators," *Nucl. Phys.* **B409** (1993) 509–546, `hep-th/9305160`.

[9] A. B. Zamolodchikov and A. B. Zamolodchikov, "Factorized S-matrices in two dimensions as the exact solutions of certain relativistic quantum field models," *Annals Phys.* **120** (1979) 253–291.

[10] D. Bernard and A. Leclair, "The quantum double in integrable quantum field theory," *Nucl. Phys.* **B399** (1993) 709–748, `hep-th/9205064`.

[11] G. W. Delius, M. D. Gould, and Y.-Z. Zhang, "On the construction of trigonometric solutions of the Yang-Baxter equation," *Nucl. Phys.* **B432** (1994) 377–403, `hep-th/9405030`.

[12] T. J. Hollowood, "Quantizing SL(N) solitons and the Hecke algebra," *Int. J. Mod. Phys.* **A8** (1993) 947–982, `hep-th/9203076`.

[13] G. M. Gandenberger and N. J. MacKay, "Exact S matrices for $d_{n+1}^{(2)}$ affine Toda solitons and their bound states," *Nucl. Phys.* **B457** (1995) 240–272, `hep-th/9506169`.

[14] G. M. Gandenberger, N. J. MacKay, and G. M. T. Watts, "Twisted Algebra R-Matrices and S-Matrices for $b_n^{(1)}$ Affine Toda Solitons and their Bound States," *Nucl. Phys.* **B465** (1996) 329–349, `hep-th/9509007`.

[15] P. Bowcock, E. Corrigan, P. E. Dorey, and R. H. Rietdijk, "Classically integrable boundary conditions for affine Toda field theories," *Nucl. Phys.* **B445** (1995) 469–500, `hep-th/9501098`.

[16] P. Bowcock, E. Corrigan, and R. H. Rietdijk, "Background field boundary conditions for affine Toda field theories," *Nucl. Phys.* **B465** (1996) 350–364, `hep-th/9510071`.

[17] G. W. Delius, "Soliton-preserving boundary condition in affine Toda field theories," *Phys. Lett.* **B444** (1998) 217–223, `hep-th/9809140`.

[18] G. W. Delius, "Restricting affine Toda theory to the half-line," *JHEP* **09** (1998) 016, `hep-th/9807189`.

[19] I. V. Cherednik, "Factorizing particles on a half line and root systems," *Theor. Math. Phys.* **61** (1984) 977–983.

[20] S. Ghoshal and A. B. Zamolodchikov, "Boundary S matrix and boundary state in two-dimensional integrable quantum field theory," *Int. J. Mod. Phys.* **A9** (1994) 3841–3886, `hep-th/9306002`.

[21] G. M. Gandenberger, "New non-diagonal solutions to the $a_n^{(1)}$ boundary Yang- Baxter equation," `hep-th/9911178`.

[22] A. Lima-Santos, "Reflection K-Matrices for 19-Vertex Models," *Nucl. Phys.* **B 558** (1999) 637–667, `solv-int/9906003`.

[23] C.-x. Liu, G.-x. Ju, S.-k. Wang, and K. Wu, "Classification of Solutions to Reflection Equation of Two- Component Systems," *J. Phys.* **A32** (1999) 3505–3523, `hep-th/9808083`.

[24] H. J. de Vega and A. Gonzalez-Ruiz, "Exact Bethe Ansatz solution for $A(n-1)$ chains with non$SU_q(n)$ invariant open boundary conditions," `hep-th/9404141`.

[25] M. T. Batchelor, V. Fridkin, A. Kuniba, and Y. K. Zhou, "Solutions of the reflection equation for face and vertex models associated with $A_n^{(1)}, B_n^{(1)}, C_n^{(1)}, D_n^{(1)}$ and $A_n^{(2)}$," *Phys. Lett.* **B376** (1996) 266–274, `hep-th/9601051`.

[26] C. Ahn and C.-K. You, "Complete non-diagonal reflection matrices of RSOS/SOS and hard hexagon models," *J. Phys. A: Math. Gen.* **31** (1998), no. 9, 2109–2121.

[27] A. Lima-Santos, "$D_{n+1}^{(2)}$ reflection K-matrices," *Nucl. Phys.* **B612** (2001), no. 3, 446–460.

[28] R. E. Behrend and P. A. Pearce, "A construction of solutions to reflection equations for interaction-round-a-face models," *J. Phys. A: Math. Gen.* **29** (1996) 7827–7836, `hep-th/9512218`.

[29] A. Doikou and P. P. Martin, "Hecke algebraic approach to the reflection equation for spin chains," `hep-th/0206076`.

[30] G. W. Delius and N. J. MacKay, "Quantum group symmetry in sine-Gordon and affine Toda field theories on the half-time," `hep-th/0112023`.

[31] G. W. Delius and R. I. Nepomechie, "Solutions of the boundary Yang-Baxter equation for arbitrary spin," *J. Phys. A: Math. Gen.* **35** (April, 2002) L341–L348, `hep-th/0204076`.

[32] E. K. Sklyanin, "Boundary conditions for integrable quantum systems," *J. Phys.* **A21** (1988) 2375.

[33] G. W. Delius, N. J. MacKay, and B. J. Short, "Boundary remnant of Yangian symmetry and the structure of rational reflection matrices," *Phys. Lett.* **B522** (2001) 335–344, `hep-th/0109115`.

[34] N. J. MacKay, "Rational K-matrices and representations of twisted Yangians," `math.qa/0205155`.

[35] P. Baseilhac and K. Koizumi, "Sine-Gordon quantum field theory on the half-line with quantum boundary degrees of freedom", `hep-th/0208005`.

[36] G. W. Delius and A. George, "Quantum affine reflection algebras of type $d_n^{(1)}$ and reflection matrices", *submitted to Lett. Math. Phys.* (2002) `math.QA/0208043`.

[37] A. I. Molev, E. Ragoucy, P. Sorba, "Coideal subalgebras in quantum affine algebras", `math.QA/0208140`.

Department of Mathematics, University of York, York YO10 5DD, United Kingdom
E-mail address: `gwd2@york.ac.uk`
URL: `http://www.york.ac.uk/mathematics/physics/delius/`